NSP Snapshots Study

COHEALTH BRAYBROOK

Results from the third quarterly two-week snapshot: September 2020 The NSP Snapshots study aims to identify service use trends and describe service user experiences, needs and outcomes of Victorian Needle and Syringe

Programs. This snapshot ran from 31 Aug – 13 Sept 2020.






contacts



contacts

Since Stage 4 COVID-19 restrictions began,

29% use decreased



Of which: 88% were unable to afford their drug of choice

46% of contacts reported that their drug use remained stable

25% use increased



4070



of contacts had experienced interactions with police



think drugs are harder to access (June survey)

Last drug injected:



Sept 2020:

June 2020:















21%

19%

Other: cocaine, MDMA, mixtures | PIEDs: Performance or Image Enhancing Drugs







NSP Snapshots Study

HEALTHWORKS FOOTSCRAY

Results from the third quarterly two-week snapshot: September 2020 CONECTA



The NSP Snapshots study aims to identify service use trends and describe service user experiences, needs and outcomes of Victorian Needle and Syringe Programs. This snapshot ran from 31 Aug – 13 Sept 2020.



64 contacts




59 contacts



21 contacts

Since Stage 4 COVID-19 restrictions began,


51% use decreased



Of which: 45% were unable to afford their drug of choice

37% of contacts reported that their drug use remained stable

13% use increased




19%



of contacts had experienced interactions with police

36%




think drugs are **harder** to access (June survey)

32%



had been tested for COVID-19 (June survey)

Last drug injected:



Sept 2020:

June 2020:

Heroin Methlores









63%

27%

<5

<5

<5

71%

14%

/[

<5

Other: cocaine, MDMA, mixtures | PIEDs: Performance or Image Enhancing Drugs







NSP Snapshots Study COHEALTH FOOTSCRAY & BRAYBROOK

Results from the third quarterly two-week snapshot: September 2020

The NSP Snapshots study aims to identify service use trends and describe service user experiences, needs and outcomes of Victorian Needle and Syringe Programs. This snapshot ran from 31 Aug — 13 Sept 2020.



FOOTSCRAY



64 contacts BRAYBROOK



56 contacts


Since Stage 4 COVID-19 restrictions began,

51% use decreased



Of which: 45% were unable to afford their drug of choice

29% use decreased



Of which: 88% were unable to afford their drug of choice

37% of contacts reported that their drug use remained stable

13% use increased



46% of contacts reported that their drug use remained stable

25% use increased

1990



of contacts had experienced interactions with police

Of which:

8 (7%) nothing happened

6 (5%) were issued a warning

7 (6%) were issued a fine

20%



Last drug injected:




Footscray:

Braybrook:

















21%

Other: cocaine, MDMA, mixtures | PIEDs: Performance or Image Enhancing Drugs





